

A LIOUVILLE PROPERTY OF HOLOMORPHIC MAPS

CHENGJIE YU

ABSTRACT. In this article, we prove a Liouville property of holomorphic maps from a complete Kähler manifold with nonnegative holomorphic bisectional curvature to a complete simply connected Kähler manifold with a certain assumption on the sectional curvature.

Liouville property is an interesting topic in analysis since Liouville found that a bounded holomorphic function on the complex plane must be a constant function. It has been studied by many geometers. For example, Yau [5] studied Liouville property of harmonic functions on complete Riemannian manifold with nonnegative Ricci curvature and Tam [4] studied Liouville property of harmonic maps. By Yau's Schwartz lemma([6]), every holomorphic map from a complete Kähler manifold with nonnegative Ricci curvature to a complete Kähler manifold with holomorphic bisectional curvature not greater than a negative constant is a constant map. This is a Liouville property of holomorphic maps.

In this article, we prove the following Liouville property of holomorphic maps.

Theorem 0.1. *Let M be a complete Kähler manifold with nonnegative bisectional curvature and N be a complete simply connected Kähler manifold with sectional curvature $\leq -\frac{c}{(1+r)^2}$ where c is some positive constant and r is the distance function of N with respect to a fixed point. Then, any holomorphic map from M to N must be constant.*

In order to prove the theorem, we need the following lemma on the existence of plenty of bounded strictly plurisubharmonic functions so that they separate points on the manifold.

Lemma 0.1. *Let M be a complete simply connected Kähler manifold with sectional curvature $\leq -\frac{c}{(1+r)^2}$ where c is some positive constant and r is the distance function of M with respect to a fixed point. Then,*

Date: Jan 2010.

2000 Mathematics Subject Classification. Primary 53B25; Secondary 53C40.

Key words and phrases. Liouville property, Kähler manifolds, negative sectional curvature.

for any point $p \in M$, there is a bounded continuous strictly plurisubharmonic function ϕ such that $\phi > 0$ on $M \setminus \{p\}$ and $\phi(p) = 0$.

Proof. We construct the strictly plurisubharmonic function with the form $\phi = f(r)$ where r is the distance function to p and f is a function to be determined.

By Hessian comparison, we have

$$(0.1) \quad D^2 r \geq \frac{\sqrt{c}}{1+r} \coth \left(\frac{\sqrt{c}r}{1+r} \right) (g - dr \otimes dr).$$

Let $e_1 = \frac{1}{\sqrt{2}}(\nabla r - J\nabla r)$ and $e_1, e_2, \dots, e_n$ be a parallel unitary frame along geodesic rays emanating from p . Then

$$(0.2) \quad r_1 = \frac{1}{\sqrt{2}}$$

and $r_\alpha = 0$ for any $\alpha > 1$.

By (0.1), when assuming $f' > 0$, we have

$$(0.3) \quad \begin{aligned} \phi_{\alpha\bar{\beta}} &= f'(r)r_{\alpha\bar{\beta}} + f''(r)r_\alpha r_{\bar{\beta}} \\ &\geq f'(r)\frac{\sqrt{c}}{1+r} \coth \left(\frac{\sqrt{c}r}{1+r} \right) (g_{\alpha\bar{\beta}} - r_\alpha r_{\bar{\beta}}) \\ &\quad + f''(r)r_\alpha r_{\bar{\beta}} \\ &:= \psi_{\alpha\bar{\beta}} \end{aligned}$$

Note that when $\alpha \neq \beta$, we have $\psi_{\alpha\bar{\beta}} = 0$, when $\alpha > 1$,

$$(0.4) \quad \psi_{\alpha\bar{\alpha}} = \frac{\sqrt{c}}{1+r} \coth \left(\frac{\sqrt{c}r}{1+r} \right) f'(r),$$

and

$$(0.5) \quad \psi_{1\bar{1}} = \frac{1}{2} \left[\frac{\sqrt{c}}{1+r} \coth \left(\frac{\sqrt{c}r}{1+r} \right) f'(r) + f''(r) \right]$$

So, if $f'(r) > 0$ and

$$(0.6) \quad [\log f']'(r) > -\frac{\sqrt{c}}{1+r} \coth \left(\frac{\sqrt{c}r}{1+r} \right),$$

then ϕ is strictly plurisubharmonic. Note that

$$(0.7) \quad \sqrt{c} \coth \left(\frac{\sqrt{c}r}{1+r} \right) \rightarrow \sqrt{c} \coth \sqrt{c}$$

as $r \rightarrow \infty$ and $\sqrt{c} \coth \sqrt{c} > 1$. So, there are two positive numbers R and δ , such that

$$(0.8) \quad \frac{\sqrt{c}}{1+r} \coth \left(\frac{\sqrt{c}r}{1+r} \right) > \frac{1+\delta}{1+r}$$

for any $r \geq R$.

Let $h(r)$ be a continuous function on $[0, \infty)$ such that

$$(0.9) \quad \begin{cases} h(r) = \frac{1+\delta}{1+r} < \frac{\sqrt{c}}{1+r} \coth\left(\frac{\sqrt{c}r}{1+r}\right) & r \geq R \\ h(r) < \frac{\sqrt{c}}{1+r} \coth\left(\frac{\sqrt{c}r}{1+r}\right) & \text{for } r \leq R \end{cases}$$

Then

$$(0.10) \quad f(r) = \int_0^r e^{-\int_0^s h(t)dt} ds$$

satisfies (0.6) and $f' > 0$. So $\phi = f(r)$ is a strictly plurisubharmonic function. Moreover,

$$(0.11) \quad \begin{aligned} f(r) &= \int_0^r e^{-\int_0^R h(t)dt - \int_R^s h(t)dt} ds \\ &= \int_0^r e^{-\int_0^R h(t)dt - \int_R^s \frac{1+\delta}{1+t} dt} ds \\ &= (1+R)^{1+\delta} e^{-\int_0^R h(t)dt} \int_0^s \frac{1}{(1+s)^{1+\delta}} ds \\ &\leq (1+R)^{1+\delta} e^{-\int_0^R h(t)dt} \int_0^\infty \frac{1}{(1+s)^{1+\delta}} ds. \end{aligned}$$

So, f is bounded. It is clear that $f(0) = 0$ and $f(r) > 0$ for any $r > 0$. Therefore, $\phi = f(r)$ is a bounded continuous strictly plurisubharmonic function on M with $\phi > 0$ on $M \setminus \{p\}$ and $\phi(p) = 0$. $\square$

Remark 0.1. Similar results can be found in [1]. Our treatment here is different with that in [1].

Proof of the theorem. We proceed by contradiction. Let $f : M \rightarrow N$ be a nonconstant holomorphic map. Then, there are two points p, q in M , such that $f(p) \neq f(q)$. By the lemma above, there is a continuous plurisubharmonic function ϕ on N such that $\phi(f(p)) = 0$, $\phi(f(q)) > 0$. Then, $\phi(f)$ is a continuous bounded plurisubharmonic function on M that is nonconstant. This contradicts Theorem 3.2 in [2]. $\square$

Remark 0.2. The unitary invariant metric on $\mathbb{C}^n$ constructed by Seshadri [3] has negative sectional curvature and sectional curvature $\leq -\frac{1}{r^2 \log r}$ outside a compact subset. This means that the curvature decay rate can not be raised to be greater than 2.

REFERENCES

- [1] Greene, R. E.; Wu, H. *Function theory on manifolds which possess a pole*. Lecture Notes in Mathematics, 699. Springer, Berlin, 1979. ii+215 pp. ISBN: 3-540-09108-4.
- [2] Ni, Lei; Tam, Luen-Fai. *Plurisubharmonic functions and the structure of complete Kähler manifolds with nonnegative curvature*. J. Differential Geom. 64 (2003), no. 3, 457–524.
- [3] Seshadri, H., *Negative sectional curvature and the product complex structure*, Math. Res. Lett. **13** (2006), 495-500.
- [4] Tam, Luen-Fai. *Liouville properties of harmonic maps*. Math. Res. Lett. 2 (1995), no. 6, 719–735.
- [5] Yau, Shing Tung. *Harmonic functions on complete Riemannian manifolds*. Comm. Pure Appl. Math. 28 (1975), 201–228.
- [6] Yau, Shing Tung. *A general Schwarz lemma for Kähler manifolds*. Amer. J. Math. 100(1978), no. 1, 197–203.

DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU, GUANG-DONG, CHINA

E-mail address: cjyu@stu.edu.cn